COMMISSION REGULATION (EEC) No 1979/79

of 5 September 1979

amending, as regards their application by variety of Community produced tobacco, Regulations (EEC) No 1727/70, (EEC) No 1728/70, (EEC) No 2603/71, (EEC) No 638/74 and (EEC) No 410/76

THE COMMISSION OF THE EUROPEAN COMMUNITIES,

Having regard to the Treaty establishing the European Economic Community,

Having regard to Council Regulation (EEC) No 727/70 of 21 April 1970 on the common organization of the market in raw tobacco (1), as last amended by Regulation (EEC) No 1579/79 (2), and in particular the first subparagraph of Article 3 (3), and Articles 5 (6), 6 (10) and 7 (4) thereof,

Having regard to Council Regulation (EEC) No 1697/71 of 26 July 1971 on the financing of intervention expenditure in respect of raw tobacco (3), as amended by Regulation (EEC) No 330/74 (4), and in particular Article 6 (2) thereof,

Whereas Council Regulation (EEC) No 1581/79 of 24 July 1979 fixing for the 1979 harvest the norm prices, intervention prices and premiums granted to purchasers of leaf tobacco and the derived intervention price of baled tobacco (5), amended the list of varieties of Community produced tobaccos and the definition of their reference qualities applied since the entry into force of the common organization of the market in raw tobacco in order to bring them more into line with the current reality of this market; whereas, therefore, amendments must be made along these lines in the Regulations the provisions of which refer to the different varieties and reference qualities of tobacco namely:

— Commission Regulation (EEC) No 1727/70 of 25 August 1970 on intervention procedure for raw tobacco (6), as last amended by Regulation (EEC) No 1556/78 (7),

— Commission Regulation (EEC) No 1728/70 of 25 August 1970 fixing the scales of price increases and reductions for raw tobacco (8), as last amended by Regulation (EEC) No 784/77 (9),

— Commission Regulation (EEC) No 2603/71 of 6 December 1971 on detailed rules for the conclusion of contracts for first processing and market preparation of tobacco held by intervention agencies (10), as last amended by Regulation (EEC) No 1018/79 (11),

— Commission Regulation (EEC) No 638/74 of 20 March 1974 fixing the maximum tolerance for quantity losses resulting from the storage of raw tobacco by intervention agencies (12),

— Commission Regulation (EEC) No 410/76 of 23 February 1976 fixing the maximum permissible weight losses in connection with the supervision of the first processing and market preparation of tobacco (13);

Whereas the measures provided for in this Regulation are in accordance with the opinion of the Management Committee for Raw Tobacco,

HAS ADOPTED THIS REGULATION:

Article 1

Annexes I, II and IV to Regulation (EEC) No 1727/70 shall be replaced respectively by Annexes I, II and III to this Regulation.

Article 2

Annexes I and II to Regulation (EEC) No 1728/70 shall be replaced respectively by Annexes IV and V to this Regulation.

Article 3

The Annex to Regulation (EEC) No 2603/71 shall be replaced by Annex VI to this Regulation.

(1) OJ No L 94, 28. 4. 1970, p. 1.
(2) OJ No L 189, 27. 7. 1979, p. 1.
(3) OJ No L 175, 4. 8. 1971, p. 8.
(4) OJ No L 37, 9. 2. 1974, p. 5.
(5) OJ No L 189, 27. 7. 1979, p. 6.
(6) OJ No L 191, 27. 8. 1970, p. 5.
(7) OJ No L 184, 6. 7. 1978, p. 11.
(8) OJ No L 191, 27. 8. 1970, p. 18.
(9) OJ No L 95, 19. 4. 1977, p. 16.
(10) OJ No L 269, 8. 12. 1971, p. 11.
(11) OJ No L 127, 24. 5. 1979, p. 11.
(12) OJ No L 77, 22. 3. 1974, p. 30.
(13) OJ No L 50, 26. 2. 1976, p. 11.

Article 4

The Annex to Regulation (EEC) No 638/74 shall be replaced by Annex VII to this Regulation.

Article 5

The Annex to Regulation (EEC) No 410/76 shall be replaced by Annex VIII to this Regulation.

Article 6

This Regulation shall enter into force on the third day following its publication in the *Official Journal of the European Communities.*

It shall be applicable for the first time to raw tobacco from the 1979 harvest.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 5 September 1979.

For the Commission

Finn GUNDELACH

Vice-President

ANNEX I

QUALITY CLASSIFICATION OF LEAF TOBACCO VARIETIES

Provided for in Article 7 (1) of Regulation (EEC) No 1727/70

1. (a) BADISCHER GEUDERTHEIMER, (b) FORCHHEIMER HAVANNA II c

'Grumpen' (Primings)

Class I Leaf ripe, sound, undamaged, reddish brown to lightly mottled.

Class II Leaf ripe, sound, mottled. Leaf edge may be overripe and brown in colour.

Class III Leaf not corresponding to Classes I and II but having the qualities required for intervention.

'Sandblatt' (Lugs)

Class I Leaf ripe, sound, undamaged, brown to slightly mottled, of uniform length.

Class II Leaf ripe, sound, mottled.

Class III Leaf not corresponding to Classes I and II but having the minimum qualities required for intervention.

'Hauptgut' (Top-quality leaf)

Class I Leaf ripe, sound, undamaged, dark brown to mottled, of uniform length (1).

Class II Leaf ripe, sound and mottled.

Class III Leaf not corresponding to Classes I and II but having the minimum qualities required for intervention.

'Obergut' (Tips)

Leaf ripe, sound, dark brown to mottled, having the minimum qualitites required for intervention.

2. BADISCHER BURLEY E

'Grumpen' (Primings)

Class I Leaf ripe, sound, undamaged, fleshy, light brown to reddish brown.

Class II Leaf ripe, sound, brown to mottled. Leaf edge may be over-ripe.

Class III Leaf not corresponding to Classes I and II but having the minimum qualities for intervention.

'Sandblatt' (Lugs)

Class I Leaf ripe, sound, undamaged, fleshy, light brown to reddish brown, of uniform length.

Class II Leaf ripe, sound, firm, brown to mottled.

Class III Leaf not corresponding to Classes I and II but having the minimum qualities required for intervention.

'Hauptgut' (Top-quality leaf)

Class I Leaf ripe, sound, undamaged, fleshy, reddish brown to light brown, of uniform length (1).

Class II Leaf ripe, sound, brown to mottled.

Class III Leaf not corresponding to Classes I and II but having the minimum qualities required for intervention.

(1) Reference quality.

'Obergut' (Tips)

Leaf ripe, sound, dark brown, having the minimum qualities required for intervention.

3. VIRGIN D

Class I Leaf ripe, sound, undamaged, yellow to reddish yellow; variations in colour such as brownish to greenish yellow are allowed on up to a third of the leaf's surface (1).

Class II Leaf ripe, sound, basically yellow; variations in colour such as brownish to greenish yellow are allowed on up to two thirds of the leaf's surface.

Class III Leaf not corresponding to Classes I and II but having the minimum qualities required for intervention.

4. (a) PARAGUAY and hybrids thereof, (b) DRAGON VERT and hybrids thereof, PHILIPPIN, PETIT GRAMMONT (Flobecq), SEMOIS, APPELTERRE

Lugs, Middle leaf or Tips

Class 1 Leaf ripe, of good combustibility, lively colour, relatively undamaged. Minor defects can be tolerated.

Class 2 Leaf with minor defects as regards combustibility, colour, texture or ripeness (over- or under-ripe) (1).

Class 3 Leaf with marked damage or defects as regards colour (pale or distinctly greenish), combustibility, soundness and firmness of texture or ripeness (clearly over- or under-ripe) but having the minimum qualities required for intervention.

Additional class for individually picked lugs

Class 1 S

Leaf ripe, of good combustibility, undamaged, not very close-grained, uniformly lively in colour.

5. NIJKERK

Feuilles basses (Leaves)

Class 1 Leaf ripe, texture still sticky, fleshy, undamaged, still crush-proof and elastic, without prominent veins, good brown colour, more or less dark, lively.

Class 2 Leaf slightly sticky, with minor defects as regards damage, veins or colour.

Class 3 Leaf brittle, not sticky, with sweating damage (defects of texture and colour) caused during curing or partially damaged by hail or disease but having the minimum qualities required for intervention.

Feuilles de tête (Tips)

Class 1 Leaf of first length (exceeding 45 cm), very sticky texture, fleshy, undamaged, crush-proof and elastic, without prominent veins, well ripened as shown by a lively brown to dark brown colour.

Class 2 Either leaf of second length (not exceeding 45 cm), having the abovementioned characteristics, or leaf of first length (exceeding 45 cm), texture still sticky, fleshy, still undamaged, crush-proof, veins more or less pronounced, of any variation of colour except bottle green (1).

Class 3 Leaf of any length, with marked defects as regards ripeness or colour (bottle green) or texture (brittle, not sticky, diseased, curing defects) but having the minimum qualities required for intervention.

(1) Middle leaves (Leaves): reference quality.

6. (a) MISIONERO and hybrids thereof, (b) RIO GRANDE and hybrids thereof

1st quality Leaf fully developed, over 45 cm, not coarse-grained, properly cured, reasonably free of damage, lively to fairly lively reddish-brown colour, satisfactory combustibility.

2nd quality Leaf fully developed, over 45 cm, not coarse-grained, lively to fairly lively light, somewhat yellowish colour, reasonably firm and undamaged, reasonably good combustibility.

3rd quality Leaf over 35 cm, sound, of more or less coarse texture, dull brown in colour on a reddish or yellowish background, moderate combustibility.

4th quality Leaf over 30 cm, with marked deterioration as regards condition or firmness of texture, or colour which may be dark, more or less greenish brown, dull or as regards excessive thickness and coarseness of texture. Moderate to poor combustibility, but having the minimum qualities required for intervention.

7. BRIGHT

Category A Leaf sufficiently ripe, without curing defects, open texture, with stems (midribs) and veins not too prominent, sound, of various shades of yellow (1).

Category B Leaf of varying consistency, possibly thin or coarse, sufficiently ripe, but with slight defects of curing and of colour which may range from mottled yellow or bronze to greenish yellow, possibly damaged.

Category C Leaf of varying consistency, of close texture, with curing defects, yellow in colour, heavily mottled or verging on grey or brown ; with serious damage but having the minimum qualities required for intervention.

8. BURLEY I

Category A Leaf sufficiently ripe, without curing defects, of open possibly firm texture, stems and veins not too prominent, sound, of a more or less lively nut-brown colour (1).

Category B Leaf of varying consistency, sound, with some curing defects and damage, not uniform in colour.

Category C Leaf coarse, of close texture, with marked curing defects, damage and colour faults, but having the minimum qualities required for intervention.

9. MARYLAND

Category A Leaf sufficiently ripe with slight curing defects and very slightly bronzed, average texture with not very prominent stems and veins, sound, rather lively reddish-brown colour.

Category B Leaf varied firmness but rather light, sound, some minor defects as regards curing and damage, reddish-brown colour not uniform.

Category C Coarse leaves with marked defects as regards curing damage and colour but having the minimum qualities required for intervention.

10. (a) KENTUCKY and hybrids thereof, (b) MORO DI CORI, (c) SALENTO

Category A Leaf undamaged, of large or average size, fully ripe, of fine texture, reasonably elastic and firm, stems and veins not too prominent, without curing defects and in good condition, uniformly brown in colour.

Category B Leaf fully ripe, of firm texture, without curing defects and in good condition, brown in colour, with some damage (1).

Category C Leaf reasonably ripe, possibly light texture, not uniform in colour, with curing defects and damage and minor defects in condition, but having the minimum qualities required for intervention.

(1) Reference quality.

11. (a) NOSTRANO DEL BRENTA, (b) RESISTENTE 142, (c) GOJANO

Category A Leaf of soft texture, fairly compact, of open grain, ripe, stems and veins not too prominent, sound, well cured or properly fermented according to the traditional method, more or less uniformly brown or light brown in colour, with no more than insignificant damage.

Category B Leaf of firm or light texture, sound, well cured or properly fermented according to the traditional method of brown or even dark brown colour, with minor damage (1).

Category C Leaf not fully ripe, not uniform in colour, possibly dark brown, coarse, with minor defects as regards curing, traditional treatment and preservation and badly broken, not qualifying for the above categories but having the minimum qualities required for intervention.

12. (a) BENEVENTANO, (b) BRASILE SELVAGGIO and similar varieties

Category A Leaf well ripened, sound, of fine and crush-proof texture, with not too prominent veins, of open grain, well cured and properly fermented, uniformly brown or possibly dark brown in colour, with some damage.

Category B Leaf sufficiently ripe, firm texture or possibly coarse or thin, with some minor defects as regards curing, fermentation and damage (1).

Category C Leaf of coarse texture, dark in colour, with marked defects as regards ripeness, curing, condition and damage but having the minimum qualities required for intervention.

13. XANTI-YAKA

14. PERUSTITZA

15. ERZEGOVINA, and similar varieties

Category A Leaf ripe, sound, carefully handled, light brown or yellow in colour, rather glossy, of light or fairly firm open texture, mainly of small or medium size usually obtained from tips and higher middle leaf (varieties 13 and 14) and from tips or middle leaf (variety 15).

Category B Leaf reasonably sound and ripe with some slight handling defects, mainly light in texture ; from yellow to brown in colour with marked damage but well preserved including lugs and primings (1).

Category C Light coarse leaf with curing defects, colour not uniform, with defects as regards damage from all the stalk positions but having the minimum qualities required for intervention.

16. (a) ROUND TIP, (b) SCAFATI, (c) SUMATRA I (2)

Category A Lugs sorted by length in the following proportions :

1st length : 15 %,
2nd length : 55 %,
3rd length : 30 %.

Leaf of convenient size fully ripe, reasonably uniform in colour, sound, of satisfactory texture, open-grained, stems and veins not prominent, well preserved, good combustibility, typical taste and flavour, suitable for wrapping cigars. May include about 20 % of broken leaf.

Category B Lower middle leaf sorted by length in the following proportions :

1st length : 60 %,
2nd length : 35 %,
3rd length : 5 %.

Leaf of convenient size, fully ripe and uniform in colour, sound, without damage, fine texture, elastic and crush-proof, stems and veins not prominent, well fermented and in good condition, good combustibility, of typical flavour and aroma, suitable for wrapping cigars including about 25 % of broken leaf (1).

(1) Reference quality.
(2) Lengths are as follows :
1st length : equal to or exceeding 38 cm,
2nd length : from 32 to under 38 cm,
3rd length : from 25 to under 32 cm.

Category C Higher middle leaf sorted by length in the following proportions :

1st length : 10 %,
2nd length : 40 %,
3rd length : 50 %.

Leaf of medium size, reasonably ripe, colour reasonably uniform, fairly firm texture, in good condition, medium combustibility, typical flavour and aroma, usable to some extent for wrapping cigars of inferior quality. May include about 30 % of broken leaf. Must have minimum qualities required for intervention.

ANNEX II

QUALITY CLASSIFICATION OF BALED TOBACCO

provided for in Article 7 (2) of Regulation (EEC) No 1727/70

1. (a) BADISCHER GEUDERTHEIMER, (b) FORCHHEIMER HAVANNA II c

'Grumpen' (Primings)

Class I Leaf ripe, sound, undamaged, slightly mottled to dark brown in colour; normally fermented.

Class II Leaf ripe, sound, basic colour brownish, differences in colour are accepted; the edges of the leaves may be over-ripe; normally fermented.

Class III Leaf not corresponding to Classes I and II; normally fermented.

'Sandblatt' (Lugs)

Class I Leaf ripe, sound, undamaged, slightly mottled to dark brown in colour; of uniform length; normally fermented.

Class II Leaf ripe, sound, basic colour brownish, differences in colour are accepted; normally fermented.

Class III Leaf not corresponding to Classes I and II; normally fermented.

'Hauptgut' (Leaves)

Class I Leaf ripe, sound, undamaged, slightly mottled to dark brown in colour, uniform length; normally fermented (1).

Class II Leaf ripe, sound, basic colour brownish, differences in colour are accepted; normally fermented.

Class III Leaf not corresponding to Classes I and II; normally fermented.

'Obergut' (Tips)

Leaf ripe, sound, different colours; normally fermented.

2. BADISCHER BURLEY E

'Grumpen' (Primings)

Class I Leaf ripe, sound, undamaged, fleshy, bright brown, reddish brown to brown in colour; normally fermented.

Class II Leaf ripe, sound, basic colour brownish, differences in colour are accepted; the edges of the leaf may be over-ripe; normally fermented.

Class III Leaf not corresponding to Classes I and II; normally fermented.

'Sandblatt' (Lugs)

Class I Leaf ripe, sound, undamaged, fleshy, bright brown, reddish brown to brown in colour; of uniform length; normally fermented.

Class II Leaf ripe, sound, firm, basic colour brownish; differences in colour are accepted; normally fermented.

Class III Leaf not corresponding to Classes I and II; normally fermented.

'Hauptgut' (Leaves)

Class I Leaf ripe, sound, undamaged, fleshy, bright brown, reddish brown to dark brown in colour; of uniform length; normally fermented (1).

Class II Leaf ripe, sound, basic colour brownish; differences in colour are accepted; normally fermented.

Class III Leaf not corresponding to Classes I and II; normally fermented.

(1) Reference quality.

'Obergut' (Tips)

Leaf ripe, sound, different colours ; normally fermented.

3. VIRGIN D

Class I Leaf ripe, sound, undamaged, yellow, yellow-red to brownish yellow in colour ; normally fermented (1).

Class II Leaf ripe, sound, basic colour yellow, differences in colour are accepted ; normally fermented.

Class III Leaf not corresponding to Classes I and II ; normally fermented.

7. BRIGHT

Category A Leaf reasonably ripe, carefully handled, open texture with stems and veins not too prominent, sound, of various shades of yellow (1).

Category B Leaf of varying consistency, possibly thin or coarse, sufficiently ripe with minor defects as regards handling and colour (mottled yellow or bronzed) ; may be damaged.

Category C Leaf of varying consistency, close texture, with handling defects, but suitable for storage, heavily mottled, yellow in colour or verging on either grey or brown ; with serious damage.

8. BURLEY I

Category A Leaf reasonably ripe, carefully handled, of open possibly firm texture, with stems and veins not too prominent, sound, of more or less lively nut-brown colour (1).

Category B Leaf of varying consistency, sound, with some handling defects, some damage, not uniform in colour.

Category C Coarse leaf of close texture with marked handling and colour defects, but suitable for storage.

9. MARYLAND

Category A Leaf reasonably ripe, minor curing defects and very few bronzed leaves, however, carefully handled, average texture, stems and veins not too prominent, sound, warm reddish brown in colour.

Category B Leaf of varying consistency, rather light, sound, with some defects as regards curing and damage, reddish brown in colour, not uniform.

Category C Coarse leaves with marked defects as regards curing, damage and colour but suitable for storage.

10. (a) KENTUCKY and hybrids thereof, (b) MORO DI CORI, (c) SALENTO

Category A Leaf large or medium size, intact, fully ripe, of fine reasonably elastic and firm texture, with stems and veins not too prominent, carefully handled and preserved, uniformly brown in colour.

Category B Leaf fully ripe, of firm texture, carefully handled and preserved, brown in colour, with some damage (1).

Category C Leaf reasonably ripe, may be of light texture, not uniform in colour, with handling defects, some damage, and minor defects of condition.

11. (a) NOSTRANO DEL BRENTA, (b) RESISTENTE 142, (c) GOJANO

Category A Leaf of delicate texture, reasonably firm with open grain, ripe, with stems and veins not too prominent, sound, carefully handled and fermented, of a more or less uniform brown or light brown colour, may have minor damage.

Category B Leaf of firm or light texture, sound, carefully handled and fermented, brown or dark brown in colour, with not too marked damage (1).

Category C Leaf not fully ripe, not uniform in colour, may be dark brown, coarse, with minor defects of handling, fermentation and condition with badly torn leaf, not qualifying for the above categories.

(1) Reference quality.

12. (a) BENEVENTANO, (b) BRASILE SELVAGGIO and similar varieties

Category A Well ripened leaf, sound, having a fine, crush-proof, open texture, with stems and veins not too prominent, carefully handled and fermented, brown or even dark brown but uniform in colour, with damage.

Category B Leaf reasonably ripe, of firm or even coarse or thin texture, with handling and fermentation defects and minor damage (1).

Category C Leaf of coarse texture, with marked defects as regards ripeness, handling and condition and marked damage.

13. XANTI-YAKA

14. PERUSTITZA

15. ERZEGOVINA and similar varieties

Category A Leaf ripe, healthy, carefully handled, light brown or yellow in colour, rather glossy, delicate or averagely consistent texture, open texture or small or medium size, normally from the upper or higher middle leaf (varieties 13 and 14) and the top or middle leaf (variety 15).

Category B Reasonably sound and ripe, some minor curing defects, texture most often light, colour going from yellow to red with marked defects as regards damage but in good condition, even from the lugs (1).

Category C Light and coarse leaf, with defects as regards curing but capable of being stored, colour not uniform, marked damage from all the stalk positions.

16. (a) ROUND TIP, (b) SCAFATI, (c) SUMATRA I (2)

Category A Lugs and primings, sorted by length in the following proportions :

1st length : 15 %,
2nd length : 55 %,
3rd length : 30 %.

Leaf of convenient size, fully ripe, reasonably uniform in colour, sound, of satisfactory texture, open grain, stems and veins not prominent, in good condition, good combustibility, of typical flavour and aroma, suitable for wrapping cigars. May include about 20 % broken leaf.

Category B Lower middle leaf, sorted by length in the following proportions :

1st length : 60 %,
2nd length : 35 %,
3rd length : 5 %.

Leaf of convenient size, fully ripe and uniform in colour, sound without damage, of a fine elastic and crush-proof texture, stems and veins not prominent, well fermented and in good condition, good combustibility, of typical flavour and aroma, suitable for wrapping cigars including about 25 % broken leaf (1).

Category C Higher middle leaf, sorted by length in the following proportions :

1st length : 10 %,
2nd length : 40 %,
3rd length : 50 %.

Leaf of medium size, reasonable ripeness, and uniformity of colour, of fairly firm texture, in good condition, medium combustibility, of typical flavour and aroma, useful to some extent for wrapping cigars of inferior quality. May include about 30 % broken leaf.

(1) Reference quality.
(2) The lengths are as follows :
1st length : equal to or exceeding 38 cm,
2nd length : from 32 to under 38 cm,
3rd length : from 25 to under 32 cm.

ANNEX III

Moisture to be taken into consideration when determining the net weight referred to in Article 6 (1) of Regulation (EEC) No 1727/70

Serial No	Varieties	Moisture (%)	
		Leaf tobacco	Baled tobacco
1	(a) Badischer Geudertheimer (b) Forchheimer Havanna II c	26	16
2	Badischer Burley E	25	15
3	Virgin D	19	13
4	(a) Paraguay and hybrids thereof (b) Dragon vert and hybrids thereof Philippin Petit Grammont (Flobecq) Semois Appelterre	27	19
5	Nijkerk	27	19
6	(a) Misionero and hybrids thereof (b) Rio Grande and hybrids thereof	27	19
7	Bright	16	13
8	Burley I	19	13
9	Maryland	19	13
10	(a) Kentucky and hybrids thereof (b) Moro di Cori (c) Salento	23	16
11	(a) Nostrano del Brenta (b) Resistente 142 (c) Gojano	22	18
12	(a) Beneventano (b) Brasile Selvaggio and similar varieties	24	16
13	Xanti-Yaka	17	13
14	Perustitza	17	13
15	Erzegovina and similar varieties	17	13
16	(a) Round Tip (b) Scafati (c) Sumatra I	22	16

ANNEX IV

Scale of the prices and reductions for leaf tobacco referred to in Article 1 (1) of Regulation (EEC) No 1728/70

Serial No	Varieties	Stalk position	Classes, quality and number indices in relation to reference qualities			
			Class I	Class II	Class III	
1	(a) Badischer Geudertheimer (b) Forchheimer Havanna II c	Grumpen (Primings) Sandblatt (Lugs) Hauptgut (Leaves) Obergut (Tips)	90 109 100 (1) 67	73 88 82	58 71 70	
			Class I	Class II	Class III	
2	Badischer Burley E	Grumpen (Primings) Sandblatt (Lugs) Hauptgut (Leaves) Obergut (Tips)	114 122 100 (1) 50	89 98 77	56 76 59	
			Class I	Class II	Class III	
3	Virgin D		100 (1)	57	22	
			Class 1 S	Class 1	Class 2	Class 3
4	(a) Paraguay and hybrids thereof (b) Dragon vert and hybrids thereof Philippin, Petit Grammont (Flobecq), Semois, Appleterre	Lugs Middle Leaves (Leaves) Top Leaves (Tips)	130 — —	110 120 100	90 100 (1) 80	70 80 60
			Class 1	Class 2	Class 3	
5	Nijkerk	Lugs (Leaves) Top Leaves (Tips)	100 120	80 100 (1)	60 60	
			1st quality	2nd quality	3rd quality	4th quality
6	(a) Misionero and hybrids thereof (b) Rio Grande and hybrids thereof		120	100 (1)	80	60
			Category A	Category B	Category C	
7	Bright		100 (1)	65	50	
8	Burley I		100 (1)	65	48	
9	Maryland		100 (1)	65	48	
10	(a) Kentucky and hybrids thereof (b) Moro di Cori (c) Salento		157	100 (1)	60	
11	(a) Nostrano del Brenta (b) Resistente 142 (c) Gojano		129	100 (1)	76	
12	(a) Beneventano (b) Brasile Selvaggio and similar varieties		128	100 (1)	71	
13	Xanti-Yaka		116	100 (1)	74	
14	Perustitza		109	100 (1)	73	
15	Erzegovina and similar varieties		109	100 (1)	71	
16	(a) Round Tip (b) Scafati (b) Sumatra I		86	100 (1)	36	

(1) Reference quality.

ANNEX V

Scales of price increases and reductions for baled tobacco referred to in Article 1 (2) of Regulation (EEC) No 1728/70

Price increases and reductions in relation to classification categories

Serial No	Varieties	Stalk position	Classes and number indices in relation to reference qualities		
			Class I	Class II	Class III
1	(a) Badischer Geudertheimer (b) Forchheimer Havanna II c	Grumpen (Primings) Sandblatt (Lugs) Hauptgut (Leaves) Obergut (Tips)	90 109 100 (1) 67	73 88 82	58 71 70
2	Badischer Burley E	Grumpen (Primings) Sandblatt (Lugs) Hauptgut (Leaves) Obergut (Tips)	114 122 100 (1) 50	89 98 77	56 76 59
3	Virgin D		100 (1)	57	22
			Category A	Category B	Category C
7	Bright		100 (1)	65	50
8	Burley I		100 (1)	65	48
9	Maryland		100 (1)	65	48
10	(a) Kentucky and hybrids thereof (b) Moro di Cori (c) Salento		157	100 (1)	60
11	(a) Nostrano del Brenta (b) Resistente 142 (c) Gojano		129	100 (1)	76
12	(a) Beneventano (b) Brasile Selvaggio and similar varieties		128	100 (1)	71
13	Xanti-Yaka		116	100 (1)	74
14	Perustitza		109	100 (1)	73
15	Erzegovina and similar varieties		109	100 (1)	71
16	(a) Round Tip (b) Scafati (c) Sumatra		86	100 (1)	36

(1) Reference quality.

ANNEX VI

Amount referred to in Article 6 (1) of Regulation (EEC) No 327/71

Serial No	Variety	Amount in ECU/kg of leaf tobacco
1	(a) Badischer Geudertheimer (b) Forchheimer Havanna II c	0·580
2	Badischer Burley E	0·598
3	Virgin D	0·603
4	(a) Paraguay and hybrids thereof (b) Dragon vert and hybrids thereof Philippin Petit Grammont (Flobecq) Semois Appelterre	0·422
5	Nijkerk	0·372
6	(a) Misionero and hybrids thereof (b) Rio Grande and hybrids thereof	0·468
7	Bright	0·642
8	Burley I	0·594
9	Maryland	0·594
10	(a) Kentucky and hybrids thereof (b) Moro di Cori (c) Salento	0·490
11	(a) Nostrano del Brenta (b) Resistente 142 (c) Gojano	0·589
12	(a) Beneventano (b) Brasile Selvaggio and similar varieties	0·445
13	Xanti-Yaka	0·957
14	Perustitza	0·663
15	Erzegovina and similar varieties	0·608
16	(a) Round Tip (b) Scafati (c) Sumatra I	3·598

ANNEX VII

Annex to Regulation (EEC) No 638/74

Tolerance limits as a percentage of the net weight

Varieties	Year of purchase	Following years
Dark air cured (1, 4, 5, 6, 11, 12) Fire cured (10) Special tobaccos (16)	1·5 %	0·7 %
Light air cured (2, 8, 9) Flue cured (3, 7) Sun cured (13, 14, 15)	1·0 %	0·6 %

ANNEX VIII

Maximum rate of weight losses referred to in Article 1 of Regulation (EEC) No 410/76

Serial No	Varieties	Maximum weight losses (as a percentage of the net weight of the leaf tobacco) (1)
1	(a) Badischer Geudertheimer (b) Forchheimer Havanna II c	22
2	Badischer Burley E	19
3	Virgin D	13
4	(a) Paraguay and hybrids thereof (b) Dragon vert and hybrids thereof Philippin Semois Appelterre	19
5	Nijkerk	19
6	(a) Misionero and hybrids thereof (b) Rio Grande and hybrids thereof	17
7	Bright	14
8	Burley I	15
9	Maryland	15
10	(a) Kentucky and hybrids thereof (b) Moro di Cori (c) Salento	15
11	(a) Nostrano and hybrids thereof (b) Resistente 142 (c) Gojano	15
12	(a) Beneventano (b) Brasile Selvaggio and similar varieties	15
13	Xanti-Yaka	17
14	Perustitza	17
15	Erzegovina and similar varieties	17
16	(a) Round Tip (b) Scafati (c) Sumatra I	16

(1) These rates are increased by four points for tobacco which has been thrashed.