This text is meant purely as a documentation tool and has no legal effect. The Union's institutions do not assume any liability for its contents. The authentic versions of the relevant acts, including their preambles, are those published in the Official Journal of the European Union and available in EUR-Lex. Those official texts are directly accessible through the links embedded in this document

►**B** **COMMISSION IMPLEMENTING REGULATION (EU) No 837/2012**

of 18 September 2012

concerning the authorisation of 6-phytase (EC 3.1.3.26) produced by *Aspergillus oryzae* (DSM 22594) as feed additive for poultry, weaned piglets, pigs for fattening and sows (holder of authorisation DSM Nutritional Products)

(Text with EEA relevance)

(OJ L 252, 19.9.2012, p. 7)

Amended by:

		Official Journal		
		No	page	date
►**M1**	Commission Implementing Regulation (EU) No 1265/2012 of 17 December 2012	L 356	61	22.12.2012
►**M2**	Commission Implementing Regulation (EU) 2016/1881 of 24 October 2016	L 289	15	25.10.2016

▼B

COMMISSION IMPLEMENTING REGULATION (EU) No 837/2012

of 18 September 2012

concerning the authorisation of 6-phytase (EC 3.1.3.26) produced by *Aspergillus oryzae* (DSM 22594) as feed additive for poultry, weaned piglets, pigs for fattening and sows (holder of authorisation DSM Nutritional Products)

(Text with EEA relevance)

Article 1

The preparation specified in the Annex, belonging to the additive category ‘zootechnical additives’ and to the functional group ‘digestibility enhancers’, is authorised as an additive in animal nutrition, subject to the conditions laid down in that Annex.

Article 2

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union.*

This Regulation shall be binding in its entirety and directly applicable in all Member States.

▼M2

ANNEX

Identification number of the additive	Name of the holder of authorisation	Additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
						Units of activity/kg of complete feedingstuff with a moisture content of 12 %			
Category of zootechnical additives. Functional group: digestibility enhancers.									
4a18	DSM Nutritional Products Ltd	6-phytase (EC 3.1.3.26)	*Additive composition* Preparation of 6-phytase (EC 3.1.3.26) produced by *Aspergillus oryzae* (DSM 22594) with a minimum activity of: 10 000 FYT (1)/g in solid form 20 000 FYT/g in liquid form *Characterisation of the active substance* 6-phytase (EC 3.1.3.26) produced by *Aspergillus oryzae* (DSM 22594) *Analytical method* (2) For quantification of 6-phytase in feed: Colorimetric method measuring the inorganic phosphate released by the 6-phytase from phytate (ISO 30024:2009)	Poultry Pigs for fattening Piglets (weaned)	—	500 FYT	—	1. In the directions for use of the additive and premixture, indicate the storage temperature, storage life, and stability to pelleting. 2. Recommended dose per kilogram of complete feed for: — poultry, piglets (weaned) and pigs for fattening: 500-4 000 FYT, — sows: 500-4 000 FYT. 3. For use in feed containing more than 0,23 % phytin-bound phosphorus. 4. For users of the additive and premixtures, feed business operators shall establish operational procedures and organisational measures to address potential risks resulting from its use. Where those risks cannot be eliminated or reduced to a minimum by such procedures and measures, appropriate personal protective equipment should be used.	9 October 2022
				Sows		500 FYT			

▼M2

Identification number of the additive	Name of the holder of authorisation	Additive	Composition, chemical formula, description, analytical method	Species or category of animal	Maximum age	Minimum content	Maximum content	Other provisions	End of period of authorisation
						Units of activity/kg of complete feedingstuff with a moisture content of 12 %			
								5. For use in weaned piglets up to 35 kg.	

([1]) 1 FYT is the amount of enzyme which liberates 1 μmol of inorganic phosphate from phytate per minute under reaction conditions with a phytate concentration of 5,0 mM at pH 5,5 and 37 °C.
([2]) Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx